

Attentional resolution

Sheng He, Patrick Cavanagh and James Intriligator

Attention can enhance selectively the visual information processing of particular locations or objects. Recent studies have shown that this enhancement has limited spatial resolution, the smallest regions that can be isolated by attention are much coarser than the smallest details that can be resolved by vision. Multiple similar objects spaced more finely than the limit of attentional resolution cannot be individuated for further processing and can only be perceived as a grouped texture. As a result, at any given time, only part of the spatial and temporal information registered by the early sensory systems is available to conscious perception. It is likely that attentional resolution is limited at a stage beyond V1 and that it has a finer grain in the lower visual field than in the upper field. The spatial aperture of attention is elongated along the radial axis relative to fixation. The briefest temporal window of attention is also much broader than visual temporal resolution. Many perceptual phenomena related to rapid serial visual presentation may reflect the limited temporal resolution of attention.

We have all experienced paying attention to, or concentrating on a piece of music, a book or a thought and losing awareness of other surrounding events. Attention is such a common term in everyday life that often we do not realize the variety of attentional phenomena underlying our personal experiences. Despite William James's famous claim that: 'everybody knows what attention is', we have only a preliminary notion of what mechanisms mediate the effects of attention. Many studies in the attention literature are concerned with what it is, some are concerned with how it works and some are concerned with what happens to information that is not being attended¹⁻³. In this short review, we will look at a specific question: how sharply can we focus our attention?

Many authors have used a 'spotlight' metaphor⁴⁻⁶ to convey the idea that attention can be restricted to a small area (illuminating only a small part of our visual field), and that this beam of attention can move around. In the more extended 'zoom lens' metaphor⁷⁻⁹, the beam not only moves but also scales up or down in size. Measurements of the distribution of attention within the beam suggest a gradient of attention, and even a tear-drop shape, oriented along radial lines from fixation¹⁰⁻¹⁴.

The spatial specificity and the unitary nature of the attentional focus have been challenged recently. A number of experiments have shown that attention can be directed to non-contiguous locations^{15,16} and even used to track as many as four or five randomly moving items simultaneously¹⁷. At the same time, other studies have demonstrated that attention is attached to objects, not locations^{18,19}. When two objects are superimposed but attention is directed to only one of them, attentional facilitation is found to be specific to the object and not its location.

The grain of attention

How finely can attention be focused? Imagine there are two dots presented above a fixation point. An observer can attend to one or the other easily when the two dots are far apart (no eye movements allowed). As the two dots get closer and closer, however, it becomes increasingly difficult to focus attention on a single dot to the exclusion of the other (that is, to individuate it). As we shall see, the price of being too close is to lose access to the features of an individual item – which seem to get mixed up with those of its neighbors (Box 1). Surprisingly, this phenomenon appears to happen well before the two dots start to fuse visually and appear as a single dot.


Eriksen and his colleagues⁴ proposed that the attention spotlight has a size of 1 degree, which means that if two items are spaced less than this distance apart they can no longer be accessed individually by attention. Although this number (1 degree) is valid only for the specific paradigm and stimulus placements that they used, it does suggest that attentional selection has a coarser grain than visual resolution. The limited resolution of attention was also evident in a study by Usai, Umiltà and Nicoletti²⁰. They showed that attention to a pac-man figure provided facilitation that was limited to targets within the pac-man except if the gap (the open mouth of the pac-man) was small. In that case, facilitation was also seen for targets in the gap as if the area through which attention had spread was a blurred version of the pac-man, blurred to the extent that the narrower gaps were filled in.

Since we will compare attentional resolution to visual resolution, let us begin with a simple description of visual resolution – the finest spacing at which visual detail can still be seen. Visual resolution is conventionally measured as the


*S. He, P. Cavanagh
and J. Intriligator are
in the Vision Sciences
Laboratory,
Psychology
Department,
Harvard University,
33 Kirkland Street,
Cambridge,
MA 02138, USA.*

tel: +1 617 495 3884
fax: +1 617 495 3764
e-mail: sheng@wjh.
harvard.edu

Box 1. How fine is the resolution of attention?



While fixating the cross in the center of the left-hand diagram (shown above) notice that it is fairly easy to voluntarily attend any item in the four concentric arrays. This is possible because each item is spaced at just less than the critical local density for individu-



ation. The diagram on the right has a density that exceeds the resolution limit of our attention. Consequently, while fixating the right-hand cross, it is difficult to move your attention from one item to another. (Adapted from Intriligator and Cavanagh, ARVO, 1997.)

finest sinusoidal grating that can be seen when presented at maximum contrast²¹. Under bright illumination, an observer can normally resolve up to 55 cycles per degree (imagine 110 black and white stripes painted on your index fingernail, held out at arm's length). In this context, resolve means either that the observer can tell the grating patch apart from a uniform field (detection task) or can tell its orientation (discrimination task). This resolution limit drops dramatically as the test is moved away from the center of gaze.

We can define attentional resolution in a similar manner: the finest spacing of the bars of a high-contrast grating that allows the observer to isolate or index each bar with attention. A simple test of this ability is to count the bars without eye movements. Figure 1 demonstrates the difference between conventional visual resolution and the attentional

resolution. We can see the gratings clearly and report that there are several fine bars vertically oriented. However, while fixating the central dot, it is much more difficult to individuate and count the bars in the grating to the left. We resolve the grating as textures but we cannot access the individual elements. Individuating the bars only becomes possible when the grating is much coarser, as seen on the right.

Studies on lateral inhibition or crowding

In the grating above, the cost of not being able to individuate the bars is relatively small – we do not know how many bars there are but we know that they are all vertical and thin. If each bar were subtly different, however, the cost might include the inability to report the special features of individual bars. This effect has been addressed in the extensive

literature on crowding, tested typically with multiple letters presented in a row, either as a word or a non-word letter string^{22,23}. These studies found that the letters in the middle of the row are more difficult to report than letters close to the fixation point, and surprisingly they are also more difficult to report than letters at the outermost position, even with prolonged viewing time²⁴. Clearly, eccentricity is not a sufficient explanation for this result. Some researchers have tried to explain these findings by appealing to a sensory level lateral inhibition process, ruling out attentional factors²⁵ or spatial uncertainty²⁶. Conversely, some authors have suggested that attention is a key factor^{27,28}.




Fig. 1 An illustration of the difference between visual resolution and attentional resolution. Looking at the center cross, the grating on the left is easily resolved. One can tell the contrast, orientation and spacing of the stripes. But it is much harder, or even impossible, to attend to individual bars or count through them without moving your eyes from the fixation cross. The grating on the right side, however, is much more sparse, and it is possible to individuate the bars. Consequently, we say that the grating on the left is beyond attentional resolution, and that the one on the right is resolvable by attention.

The lateral interference effect measured in these experiments is not limited to letter stimuli. Studies on orientation discrimination and vernier acuity with line segments also have shown interference effects in the presence of properly positioned surrounding lines^{29,30}. In general, when a target is embedded in multiple distractors, it is very difficult to perceive the target. It is our claim, outlined below, that this difficulty reflects to a large degree the limited spatial resolution of our attentional mechanism.

Interestingly, the crowding effect depends strongly on the similarity between the distractors and the target to be detected, and their relative depth^{29,31}. Also, grouping between the distractors and the target changes the crowding effect significantly³². These results indicate that these effects emerge at a fairly high level.

Crowding as a measure of attentional resolution

In severe cases of crowding, a target embedded in an array of distractors cannot be selected by attention, and cannot be consciously analyzed independently of the distractors. The suggestion that the crowding effect is due largely to insufficient spatial attentional resolution is addressed by the following experiment on orientation selective adaptation³³.

Adaptation to a grating in a particular orientation will raise the threshold for subsequently detecting a grating in that same orientation much more than a grating in the orthogonal orientation. This orientation selective adaptation has been linked to neurons at the primary visual cortex and beyond³⁴. In a recent study³³, we adapted our observers either to a single grating or a crowded grating array. In the latter (crowded) case, the adapting grating at the target position was essentially unavailable to conscious perception (see Fig. 2): the identification of its orientation was at chance level. However, in both the single and crowded adapting conditions, we found the same magnitude of orientation selective adaptation effect. Figure 2 shows the stimulus and the results. A grating that is unavailable to our conscious perception (due to crowding) is, nevertheless, as powerful as a single, perfectly visible grating in producing orientation specific adaptation effects. This suggests that crowding happens at a stage beyond the site of adaptation, which itself must be no earlier than V1 (no orientation analysis occurs at earlier levels). This result, like the earlier ones showing that an unresolvable high frequency grating can produce an orientation selective adaptation effect³⁵, is consistent with the proposal that activation of V1 neurons alone is not sufficient for visual awareness^{36,37}.

In the context of early versus late attentional selection, these results strongly support late selection. The lateral interference or crowding, which prevents the selection of the target grating, is certainly not at an early 'sensory' level as the information has been registered and processed to at least the level of area V1, if not higher. Is this high-level lateral interference a property of attention itself or the result of additional cortical filtering beyond area V1 but preceding attentional selection? Our example in Fig. 1 demonstrates the phenomenon of crowding occurring clearly at the level of selection. The fine vertical bars are seen clearly, and the observer can report any number of details about the lines: width, orientation, length and so on. And yet, as is revealed





Fig. 2 Orientation selective adaptation. Orientation selective adaptation in displays where observers could and could not perceive the orientation of the adapting grating (fourth grating away from fixation). (A) Contrast thresholds were measured under three different conditions: adapt to a single grating, adapt to the fourth grating in a linear array of five gratings and no adaptation (baseline). (B) Threshold contrast elevation after adaptation relative to baseline contrast threshold before adaptation. The difference between same adapt-test orientation and different adapt-test orientation represents the strength of orientation selective adaptation. The data show that a flanked grating (orientation not perceivable) and a single grating (orientation readily perceivable) were almost equally effective in orientation specific adaptation. (From Ref. 33.)

when attempting to count the lines without eye movements, the individual lines cannot be accessed. Classical crowding displays with adjacent letters, or the crowded grating patches used in our experiment³³, are not so simple as the set of lines shown in Fig. 1. It is possible that these more complex stimuli suffer from some high-level lateral interference that precedes selection. They must, nevertheless, all share the same selection difficulty so clearly seen in Fig. 1. It is this selection bottleneck that we are labeling the resolution limit of attention.


Upper/lower field and radial/tangential inhomogeneities

The grain of visual resolution is not uniform in the visual field. It drops off precipitously with increasing distance from the center of gaze. However, the resolution of attention



appears to have two additional inhomogeneities that are not characteristic of visual resolution. Firstly, a lower visual field advantage is found in many different tasks that require focused attention³³. For example, in our experiment where subjects had to determine whether an item in a prespecified position (flanked with distractors) is a target or not, no marked difference between the upper and lower field was seen for targets defined by simple features (line segments

with large orientation differences), but a pronounced lower field advantage was observed for targets defined by feature conjunctions (Fig. 3, from Ref. 33). The lower field advantage is also evident in an attentional tracking task (similar to that developed by Pylyshyn and his colleagues, Ref. 17), where observers were asked to track two discs among nine randomly moving discs (Fig. 4, from Ref. 33). Since the tracked discs are physically identical to the non-tracked



ones (except for a brief color change for identification at the beginning of the trial), performance in this task depends critically on the ability to use attention to track the selected targets, that sometimes move very close to the distractors.

Differences between the upper and lower visual field were also found in several other tasks involving attention³⁸. Previc proposed two attentional systems, one for extrapersonal space that favors the upper visual field, and one for peripersonal space that is directed towards the proximal lower visual field³⁸. Although this distinction is still a subject of debate, much of the evidence is consistent with a finer attentional resolution in the lower visual field. In contrast, there is very little evidence of differences in visual resolution between the upper and lower field within a large range of eccentricity as measured by contrast sensitivity using gratings³⁹. The recent finding that the lower visual field is better at perceiving subjective contours⁴⁰ may be unrelated to the different attentional resources of the two hemifields, although Gurnsey, Poirier and Gascon⁴¹ have argued that subjective contours may not be seen without attention.

The lower visual field advantage in attentional resolution may partly be due to the fact that the lower field is represented in the upper part of the primary visual cortex, which is anatomically adjacent to and projects more heavily⁴² into the occipital–parietal regions that are often linked to spatial attentional control^{43,44}.

The second inhomogeneity is that attentional resolution is also anisotropic in the radial or tangential dimension. There is significantly less interference between items when they are arranged tangentially, rather than radially, relative to the eyes' fixation^{45,46}. It seems that the shape of the smallest regions to which we can attend is ellipsoidal, elongated along the radial lines from fixation. Although the receptive fields of many visual neurons^{47–49} tend to be slightly elongated along such radial lines, this factor alone is not sufficient to explain the large difference between radial and tangential configurations in the lateral interaction which is observed behaviorally. The difference in attentional resolution may contribute to many observed radial orientation advantages including that for grating resolution⁵⁰, phase discrimination⁵¹ and motion sensitivity⁵².

Temporal resolution of attention

Under bright light, luminance fluctuation can be perceived as a flicker at frequencies as high as 50 Hz, suggesting that the initial stages of our visual system can resolve fast changes. In comparison, attentional processes appear to be much slower, and to have a coarser temporal resolution. In a direct analogy to the resolvable but not indivisible bars of the grating, demonstrated in Fig. 1, a spot flickering at slow rates can be perceived as appearing and disappearing (individual temporal events). However, at a higher rate, the spot is no longer seen as turning on and off discretely, but instead seems to be present continuously as a flickering spot (one cannot index a particular appearance or count the number of appearances of the spot). The rate (~4–6 Hz) at which the spot's phenomenal existence becomes continuous has been called the Gestalt flicker fusion rate by Van de Grind and his colleagues⁵³. In an experiment performed by

He and MacLeod⁵⁴, a grating with its contrast changed from a high to a low level or from a low to a high level could be perceived as an obvious change, but if the rate of change was higher than 4 Hz, observers could not distinguish between the two sequences of change.

Reduced performance with rapid serial visual presentation (RSVP) may also reflect the poor attentional resolution in time⁵⁵. For example, when detecting a probe that occurs between 100 ms and 450 ms after a correctly identified target in an RSVP sequence, observers show a marked decrease in their ability to detect that probe. This loss of performance has been labeled the attentional blink⁵⁶. The limited temporal resolution of attentional selection may be one of the factors that contribute to this phenomenon.

General discussions and future questions

Visual information is processed at multiple stages. We cannot be aware of everything at any one time, but only the things we are attending to or things that capture our attention. In this sense, attentional resolution is the bottleneck of our awareness and one might even argue that studying attention amounts to studying awareness. Although, in this article we have used the word attention as if it were a single process, it is not necessarily so^{57,58}. Exogenous and endogenous shifts in attention may be controlled separately⁵⁹. Moreover, the systems for shifting attention from one object to the next^{60,61} may be very different from those used to keep track of moving targets⁶².

A central goal in the study of attention is to identify the neural structures mediating attentional processes and the study of neuronal responses to crowded stimuli may have much to offer here. For example, in many cortical areas the response of a cell to its preferred stimulus is strongly reduced when the stimulus is flanked by identical stimuli^{63–66}. However, based on what has been found psychophysically^{45,46}, the signature of a cell which participates in attentional selection should be that the response is almost completely suppressed when the target and flanking stimuli are aligned radially from the foveal, whereas less suppression should be seen when target and flankers are aligned tangentially at equal eccentricity.

For many years the question of early vs. late selection has been the center of debate in studies of attention^{67–69}. It remains an unresolved question today. The adaptation experiment described earlier demonstrates an effective method for evaluating the processing of unselected features and the results indicate that the site of attentional selection is later than the site of orientation analyses. The advantage of this method lies in the degree to which stimuli are rendered inaccessible to attentional selection. Traditionally, the fate of unattended information is studied with brief, masked presentations. This masking may reduce the available signal enough to prevent accurate recognition but it is always possible that bits and pieces of the stimulus may reach awareness; these may be sufficient to prime or otherwise influence later measures⁷⁰. However, with an optimally crowded display (Fig. 2A), the target stimulus remains inaccessible to awareness, even with unlimited viewing, making the argument that unreported information is unavailable to awareness much more compelling. The same paradigm can

be extended to study the level of processing of many other properties of the crowded/inaccessible stimulus.

Finally, the pattern shown in Fig. 1 raises the question of when a pattern is no longer an arrangement of elements but a texture. The pattern on the left seemed necessarily a texture since its individual elements could not be accessed. However, the pattern on the right could be thought of as either a sparse texture or an arrangement of individual lines. Studies of texture segmentation may, therefore, also be related to attentional resolution. The factors that support segmentation may change qualitatively when the pattern density crosses the limit of attentional resolution⁷¹.

The concept of attentional resolution may help us in understanding many diverse phenomena, such as contextual effects, crowding, masking, some aspects of subliminal perception and perception in RSVP sequences. Moreover, the measurement of an individual's attentional resolution may turn out to be a more salient factor than visual resolution in predicting that individual's performance in attentionally demanding tasks such as driving, piloting airplanes or air traffic control.

Acknowledgements

Patrick Cavanagh and Sheng He were supported by NIH grants.

References

- 1 LaBerge, D.L. (1990) Attention *Psychol. Sci.* 1, 156-162
- 2 Johnston, W.A. and Dark, V.J. (1986) Selective attention *Annu. Rev. Psychol.* 37, 43-75
- 3 Desimone, R. and Duncan, J. (1995) Neural mechanisms of selective visual attention *Annu. Rev. Neurosci.* 18, 193-222
- 4 Eriksen, C.W.W. and Hoffman, J.E. (1972) Temporal and spatial characteristics of selective encoding from visual displays *Percept. Psychophys.* 12, 201-204
- 5 Jonides, J. (1980) Towards a model of the mind's eye's movement *Can. J. Psychol.* 34, 103-112
- 6 Jonides, J. (1983) Further toward a model of the mind's eye's movement *Bull. Psychonom. Soc.* 21, 247-250
- 7 Eriksen, C.W. and St James, J.D. (1986) Visual attention within and around the field of focal attention: a zoom lens model *Percept. Psychophys.* 40, 225-240
- 8 Klein, R. and McCormick, P. (1989) Covert visual orienting: hemifield-activation can be mimicked by zoom lens and midlocation placement strategies *Acta Psychol.* 70, 235-250
- 9 LaBerge, D. (1983) Spatial extent of attention to letters and words *J. Exp. Psychol. Hum. Percept. Perform.* 9, 371-379
- 10 Downing, C.J. and Pinker, S. (1985) The spatial structure of visual attention, in *Attention and Performance XI* (Posner, M.I. and Marin, O.S.M., eds), pp. 171-187, Lawrence Erlbaum Associates
- 11 Downing, C.J. (1988) Expectancy and visual-spatial attention: effects on perceptual quality *J. Exp. Psychol. Hum. Percept. Perform.* 14, 188-202
- 12 LaBerge, D. and Brown, V. (1986) Variations in size of the visual field in which targets are presented: an attentional range effect *Percept. Psychophys.* 40, 188-200
- 13 LaBerge, D. and Brown, V. (1989) Theory of attentional operations in shape identification *Psychol. Rev.* 96, 101-124
- 14 Andersen, G.J. and Kramer, A.F. (1993) Limits of focused attention in three-dimensional space *Percept. Psychophys.* 53, 658-667
- 15 Castiello, U. and Umiltà, C. (1992) Splitting focal attention *J. Exp. Psychol. Hum. Percept. Perform.* 18, 837-848
- 16 Kramer, A.F. and Hahn, S. (1995) Splitting the beam: distribution of attention over noncontiguous regions of the visual field *Psychol. Sci.* 6, 381-386
- 17 Pylyshyn, Z.W. and Storm, R.W. (1988) Tracking multiple independent targets: evidence for a parallel tracking mechanism *Spatial Vis.* 3, 179-197
- 18 Driver, J. and Baylis, G.C. (1989) Movement and visual attention: the spotlight metaphor breaks down *J. Exp. Psychol. Hum. Percept. Perform.* 15, 448-456
- 19 Duncan, J. (1984) Selective attention and the organization of visual information *J. Exp. Psychol. Gen.* 113, 501-517
- 20 Usai, M.C., Umiltà, C. and Nicoletti, R. (1995) Limits in controlling the focus of attention *Eur. J. Cogn. Psychol.* 7, 411-439
- 21 Campbell, F.W. and Gubisch, R.W. (1966) Optical quality of the human eye *J. Physiol.* 186, 558-578
- 22 Bouma, H. (1970) Interaction effects in parafoveal letter recognition *Nature* 226, 177-178
- 23 Bouma, H. (1973) Visual interference in the parafoveal recognition of initial and final letters of words *Vis. Res.* 13, 767-782
- 24 Townsend, J.T., Taylor, S.G. and Brown, D.R. (1971) Lateral masking for letters with unlimited viewing time *Percept. Psychophys.* 10, 375-378
- 25 Banks, W.P., Larson, D.W. and Prinzmetal, W. (1979) Asymmetry of visual interference *Percept. Psychophys.* 25, 447-456
- 26 Levi, D.M., Klein, S.A. and Yap, Y.L. (1987) Positional uncertainty in peripheral and amblyopic vision *Vis. Res.* 27, 581-597
- 27 Walley, R.E. and Weiden, T.D. (1973) Lateral inhibition and cognitive masking: a neuropsychological theory of attention *Psychol. Rev.* 80, 284-302
- 28 Wolford, G. and Chambers, L. (1983) Lateral masking as a function of spacing *Percept. Psychophys.* 33, 129-138
- 29 Andriessen, J.J. and Bouma, H. (1976) Eccentric vision: Adverse interactions between line segments *Vis. Res.* 16, 71-78
- 30 Westheimer, G., Shimamura, K. and McKee, S.P. (1976) Interference with line-orientation sensitivity *J. Opt. Soc. Am.* 66, 332-338
- 31 Kooi, F.L. et al. (1994) The effect of similarity and duration on spatial interaction in peripheral vision. Special issue: in honor of Bela Julesz: II. Texture and visual search *Spatial Vis.* 8, 255-279
- 32 Banks, W.P. and Prinzmetal, W. (1976) Configurational effects in visual information processing *Percept. Psychophys.* 19, 361-367
- 33 He, S., Cavanagh, P. and Intriligator, J. (1996) Attentional resolution and the locus of visual awareness *Nature* 383, 334-337
- 34 Movshon, J.A. and Lennie, P. (1979) Pattern-selective adaptation in visual cortical neurones *Nature* 278, 850-852
- 35 He, S., Smallman, H.S. and MacLeod, D.I.A. (1995) Neural and cortical limits on visual resolution *Invest. Ophthalmol. Vis. Sci. (Suppl. 36)*, 2010
- 36 Crick, F. and Koch, C. (1995) Are we aware of neural activity in primary visual cortex? *Nature* 375, 121-123
- 37 Koch, C. and Tootell, R.B. (1996) Visual psychophysics: stimulating brain but not mind *Nature* 383, 301-303
- 38 Previc, F.H. (1990) Functional specialization in the lower and upper visual fields in humans: its ecological origins and neurophysiological implications *Behav. Brain Sci.* 13, 519-575
- 39 Rovamo, J. and Virsu, V. (1979) An estimation and application of the human cortical magnification factor *Exp. Brain Res.* 37, 495-510
- 40 Rubin, N., Nakayama, K. and Shapley, R. (1996) Enhanced perception of illusory contours in the lower vs. upper visual hemifields *Science* 271, 651-653
- 41 Gurnsey, R., Poirier, F.J.A.M. and Gascon, E. (1996) There is no evidence that Kanizsa-type subjective contours can be detected in parallel *Perception* 25, 861-874
- 42 Maunsell, J.H. and Newsome, W.T. (1987) Visual processing in monkey extrastriate cortex *Annu. Rev. Neurosci.* 10, 363-401
- 43 Gazzaniga, M.S. and Ladavas, E. (1987) Disturbances in spatial attention following lesion disconnection of the right parietal lobe, in *Advances in Psychology, Neurophysiological and Neuropsychological Aspects of Spatial Neglect* (Vol. 45) (Jeannerod, M., ed.), pp. 203-213, Elsevier Science Publishing
- 44 Posner, M.I. et al. (1987) How do the parietal lobes direct covert attention? Special issue: selective visual attention *Neuropsychologia* 25, 135-145
- 45 Chambers, L. and Wolford, G. (1983) Lateral masking vertically and horizontally *Bull. Psychonom. Soc.* 21, 459-461
- 46 Toet, A. and Levi, D.M. (1992) The two-dimensional shape of spatial interaction zones in the parafovea *Vis. Res.* 32, 1349-1357
- 47 Schall, J.D., Perry, V.H. and Leventhal, A.G. (1986) Retinal ganglion cell dendritic fields in old-world monkeys are oriented radially *Brain Res.* 368, 18-23

- 48 Bauer, R. and Dow, B.M. (1989) Complementary global maps for orientation coding in upper and lower layers of the monkey's foveal striate cortex *Exp. Brain Res.* 76, 503-509
- 49 Smith, E.L.D. et al. (1990) Orientation bias of neurons in the lateral geniculate nucleus of macaque monkeys *Vis. Neurosci.* 5, 525-545
- 50 Rovamo, J. et al. (1982) Resolution of gratings oriented along and across meridians in peripheral vision *Invest. Ophthalmol. Vis. Sci.* 23, 666-670
- 51 Bennett, P.J. and Banks, M.S. (1991) The effects of contrast, spatial scale, and orientation on foveal and peripheral phase discrimination *Vis. Res.* 31, 1759-1786
- 52 Scobey, R.P. and Van Kan, P.L. (1991) A horizontal stripe of displacement sensitivity in the human visual field *Vis. Res.* 31, 99-109
- 53 Van de Grind, W.A., Grusser, O.J. and Lunkenheimer, H.U. (1973) Temporal transfer properties of the afferent visual system, in *Handbook of Sensory Physiology, Central Visual Information, A* (Vol. VII/3) (Jung, R., ed.), pp. 431-573, Springer-Verlag
- 54 He, S. and MacLeod, D.I.A. (1993) The perception of fluctuating contrast *Invest. Ophthalmol. Vis. Sci.* 34, 18
- 55 Shapiro, K.L. and Raymond, J.E. (1994) Temporal allocation of visual attention: inhibition or interference? in *Inhibitory Processes in Attention, Memory, and Language* (Dale Dagenbach, T.H.C., ed.), 151-188, Academic Press
- 56 Raymond, J.E., Shapiro, K.L. and Arnell, K.M. (1992) Temporary suppression of visual processing in an RSVP task: an attentional blink? *J. Exp. Psychol. Hum. Percept. Perform.* 18, 849-860
- 57 Allport, A. (1993) Attention and control: have we been asking the wrong questions? A critical review of twenty-five years, in *Attention and Performance 14: Synergies in Experimental Psychology, Artificial Intelligence, and Cognitive Neuroscience* (Meyer, D.E. and Kornbluth, S., eds), pp. 183-218, MIT Press
- 58 Posner, M.I. and Dehaene, S. (1994) Attentional networks *Trends Neurosci.* 17, 75-79
- 59 Theeuwes, J. (1994) Endogenous and exogenous control of visual selection *Perception* 23, 429-440
- 60 Posner, M.I. (1980) Orienting of attention *Q. J. Exp. Psychol.* 32, 3-25
- 61 Mackeben, M. and Nakayama, K. (1993) Express attentional shifts *Vis. Res.* 33, 85-90
- 62 Cavanagh, P. (1992) Attention-based motion perception *Science* 257, 1563-1565
- 63 Gilbert, C.D., Hirsch, J.A. and Wiesel, T.N. (1990) Lateral interactions in visual cortex *Cold Spring Harbor Symp. Quant. Biol.* 55, 663-677
- 64 Kapadia, M.K. et al. (1995) Improvement in visual sensitivity by changes in local context: parallel studies in human observers and in V1 of alert monkeys *Neuron* 15, 843-856
- 65 Lamme, V.A. (1995) The neurophysiology of figure-ground segregation in primary visual cortex *J. Neurosci.* 15, 1605-1615
- 66 Zipser, K., Lamme, V.A.F. and Schiller, P.H. (1996) Contextual modulation in primary visual cortex *J. Neurosci.* 16, 7376-7389
- 67 Broadbent, D.E. (1958) *Perception and Communication*, Pergamon
- 68 Deutsch, J.A. and Deutsch, D. (1963) Attention: some theoretical considerations *Psychol. Rev.* 70, 80-90
- 69 Treisman, A.M. (1960) Verbal cues, language, and meaning in selective attention *Q. J. Exp. Psychol.* 12, 242-248
- 70 Cheesman, J. and Merikle, P.M. (1984) Priming with and without awareness *Percept. Psychophys.* 36, 387-395
- 71 Durgin, F.H. (1995) Texture density adaptation and the perceived numerosity and distribution of texture *J. Exp. Psychol. Hum. Percept. Perform.* 21, 149-169

TICS Editorial Policy

Trends in Cognitive Sciences will be indispensable reading for all interested in this diverse field. The journal will carry a core of authoritative **Reviews**, written in an easily accessible style by leading authors, summarizing the exciting developments of the subjects you need to know about. Accompanying the Reviews will be a range of other types of articles. The **Comment** section will offer short updates and discussions of one or two ground-breaking and important primary papers. The authors of the primary papers will have the opportunity of replying to the Comments, thus extending the dialogues on their work. **Monitor** pieces summarize in 100-200 words a digest of recently published papers. **Meeting reports** convey the excitement of recent conferences or seminars, focussing on recent developments and the discussions surrounding them. **Opinion** articles are reviews of subjects written from a personal slant, and therefore highlight some of the controversies in cognition. **Books etcetera** will feature stimulating essay-reviews of recent publications, whether books, software, CD-ROMs, films, etcetera. And a list of books received for review will be published regularly. **Letters** stimulated by any of the articles published in *TICS* are welcome. Authors will be offered the chance to reply to any points raised.

